



Columbia Care™

Thera Ceed™

VAPORIZATION OIL 20:1
0.5 mL Cartridge

INGREDIENTS

Extracted Medicinal Cannabis, Natural Terpenes

Total Cannabinoids per 0.5mL Cartridge = 420mg
(THC 400mg, CBD 20mg)

PRODUCT FACTS

WARNINGS AND PRECAUTIONS

Medicinal cannabis products must be kept in the original container from which they were dispensed and removed from the original container only when used by the certified patient.

Keep secure at all times.

This medication may not be resold or transferred to another person.

This medication may cause drowsiness. If affected do not drive a vehicle or operate machinery.

Avoid alcohol.

Women should not consume during pregnancy or while breastfeeding except on the advice of your doctor.

If you experience any negative symptoms including, but not limited to, sore throat, headache, nausea or skin irritation, please contact your doctor immediately as some patients may experience an allergic reaction to various compounds contained within this product.

This product is for medicinal use only.

Use as directed by your doctor.

Store below 25°C. Protect from light.

KEEP OUT OF THE REACH AND SIGHT OF CHILDREN AND ALWAYS STORE IN A CHILD RESISTANT CONTAINER. Note - the cartridge and battery do not need to be stored separately.

DIRECTIONS

- Do not take more than directed
- Only use the vaporization pen/battery provided. Do not use any other vaporization or dosing device to administer this product

TO USE:

1. Attach your prescribed Columbia Care pre-filled cartridge to the battery, ensure threads are properly aligned and secure it by gently twisting in a clockwise direction
2. Seal your lips around the mouthpiece (taking care not to cover up the small air holes on the side of the cartridge)
3. The vaporizer device activates (an LED lights up) when you start to inhale through the mouthpiece and will cut off automatically after 10sec of inhalation (device will power itself off)
4. Inhale as directed for your prescribed dose
5. Hold your breath for up to 3 seconds, if possible, to maximise benefit and then exhale normally
6. Only use this product away from others who might otherwise become affected by it.

RECHARGING:

1. Light indicator: Flashes 10 times means it is time to recharge the battery
2. Remove the cartridge from the battery and store the cartridge safely in a child resistant container
3. Gently screw in the USB charging device to the end of the battery and then connect to a suitable USB power source (e.g. phone charger / laptop computer)
 - a. A white LED indicates the battery is charging
 - b. The white LED will flash 20 times and shuts off indicating the battery is fully charged.
 - c. Charge time is approximately 1 hour. Do not charge overnight or leave the device unattended while charging.
4. Once charged, remove the charger from the battery and store appropriately.


CAUTION:

- Do not expose device to liquids/moisture.
- Keep device away from flammable gasses/liquids
- Do not use in wet environments or extreme temperatures.
- Do not leave device exposed to direct sunlight for prolonged periods of time.
- Do not allow foreign metal objects to touch electrical contacts on the battery.
- Keep out of reach of children or pets.
- Do not drop, throw, or abuse device.
- Do not attempt to disassemble device. Damage and personal injury may occur.
- Battery Recycling: This product should not be treated as normal household waste. Instead it should be handed over to an applicable used battery collection point for recycling.

TERPENE PROFILE

Thera Ceed™

Effect: Calming (not scientifically proven)
Dominant Terpene: D-Limonene



CONTRAINDICATIONS

An absolute contraindication means that a circumstance, condition or substance could cause a life-threatening reaction. You should not use cannabis if you have one of these absolute contraindications:

- *Psychotic illness or a first-degree relative with schizophrenia*
- *Active unstable heart disease*
- *Allergy to cannabis or any components*

A relative contraindication means that caution should be used when circumstances exist that may put you at higher risk of complications when using cannabis. You should talk to your health care provider about your risk if you have any of these relative contraindications:

- *Use of combustible or vaporized products in a person with pulmonary disease*
- *Use in people with severe liver or kidney disease*
- *History of drug abuse or dependence, including alcohol abuse*
- *Pregnancy or breastfeeding*

Alcohol should not be consumed concurrently by a person who is using cannabis. Evidence suggests that alcohol causes faster absorption of THC, which leads to a more pronounced effect. When cannabis and alcohol are consumed together or within a short time of each other, you may experience nausea or vomiting, panic, anxiety or paranoia.

ADVERSE REACTIONS

Cannabis may produce side effects.

Cannabis side effects may include sleepiness or drowsiness. However, some people can experience the opposite effect including an increase in energy or restlessness. Other common side effects include short-term memory loss; increased appetite; mood changes; dizziness, trouble concentrating; feeling “high”; weakness, lack of coordination; anxiety, confusion; stomach pain, nausea, vomiting, diarrhoea; blushing; increase in blood pressure; increase in heart rate; decrease in blood pressure; hallucinations; paranoia; depression; dry eyes or eye redness/bloodshot eyes; or sleep problems (insomnia).

You should treat cannabis like you would any other medication that may impair the ability to perform complex tasks and avoid operating a motor vehicle or heavy machinery, performing child or elder care and making important decisions while under treatment.

If you experience a life-threatening emergency, you should call 999 or go to accident and emergency at your local hospital.

ADDICTION, ABUSE AND MISUSE

Some people, regardless of medication, may experience issues of abuse or addiction. The following indicators could be signs of an issue:

Tolerance: After prolonged or excessive use, a higher dose of drug may be needed to achieve the same level of effect.

Dependence: Characterized by the presence of withdrawal symptoms when the drug dose is reduced or discontinued.

Withdrawal: Mental and physical symptoms that occur when a dependent person reduces use or stops taking a drug; symptoms may be mild to severe.

Addiction: A substance use disorder that occurs when a person keeps using a drug and can't stop, despite negative consequences from use.

THIS PRODUCT IS INTENDED SOLELY FOR THE PATIENT FOR WHOM IT WAS PRESCRIBED AND DISPENSED AND MAY ONLY BE ADMINISTERED BY THAT PATIENT OR A DESIGNATED CAREGIVER FOR THAT PATIENT. MEDICINAL CANNABIS PRODUCTS MAY NOT BE RESOLD OR TRANSFERRED TO ANOTHER PERSON INCLUDING OTHER LICENSED MEDICINAL CANNABIS PATIENTS. KEEP THIS MEDICINE OUT OF THE REACH AND SIGHT OF CHILDREN AND PETS.

MEDICINAL CANNABIS PRODUCTS MUST BE KEPT SECURE AT ALL TIMES AND MUST BE KEPT IN THE ORIGINAL CONTAINER IN WHICH THEY WERE DISPENSED AND REMOVED FROM THE ORIGINAL CONTAINER ONLY WHEN READY FOR USE BY THE CERTIFIED PATIENT.

How to dispose of unwanted or expired medicines

Return used or expired prescription medication to a pharmacy or chemist that accepts controlled substances for safe disposal.



Manufactured for Columbia Care UK Ltd.
by IPS Pharma MS32879
41 Central Ave, West Molesey,
Surrey, KT8 2QZ