# PEDANIOS 26/1 EHD-CA



### **PEDANIOS 26/1 EHD-CA**

Medicinal Cannabis

THC/CBD: approx. 26 % THC

< 1 % CBD

Dominance: Sativa

Cultivar: Electric Honeydew (EHD)

Genetics: Girl Scout Cookies x 91

OG Krypt Melon


Origin: Canada (CA)

Unit: 10 g

### Strain description:

Hang-dried and hand-processed, Electric Honeydew (EHD) is an original San Rafael '71 variety that ensures the finest product quality and surprises with lively notes of melon, gas, and pine, along with a high THC content. This unique hybrid features dense, dark-green buds with bright flashes of orange pistils. The primary terpenes, myrcene and limonene, show analgesic, immunostimulant, and antidepressant properties.¹ In addition, this sativadominant cultivar is suitable for daytime use because to its energising, stimulating, and vitality-enhancing effects.


### Plasma level of THC:



Blood plasma levels of THC after oral administration or inhalation intake [modified from Grotenhermen 2003] <sup>2</sup>



#### Terpene profile:



#### Inhalation:

Onset: after 2 - 8 min

Peak effect: after 10 - 30 min

Duration: 2 - 6 h

Using a vaporiser is a very efficient, fast-acting, and medically recommended form of administration. The optimal temperature setting is between 175 and  $210^{\circ}\text{C}$ .

Flowers with a high THC content limit the negative effects of vaporisation, as patients can consume less to achieve the desired effect.

# PEDANIOS 26/1 EHD-CA

#### **TITRATION GUIDE**

- There is currently no uniform starting dose for cannabis products nor uniform dosing information.
- The recommended starting dose should be 0.05 g of flower. For the PEDANIOS 26/1 variety, this corresponds to approx. 13 mg of THC. Depending on effectiveness and tolerability, the dose can be increased every 1 3 days at a maximum of 0.1 g of flower.<sup>3</sup>

#### Onset and duration of effects for inhalation and oral administration<sup>4</sup>

	Inhalation		Oral administration	
Onset of Effect	<u>.</u>	Within seconds to minutes	<u>(L)</u>	Within 30 minutes to 2 hours
Peak Effects	(L)	Full effects peak within 30 minutes	Ŀ	Full effects peak within 4 hours
Duration of Effect	<u>(L)</u>	Up to 6 hours or longer	<u>(L)</u>	Up to 12 hours or longer

Adverse events should be reported to MHRA or your doctor or pharmacist.

Reporting forms and information can be found at: www.mhra.gov.uk/yellowcard



Aurora Medicine UK Limited | 5th Floor | 14-16 Dowgate Hill | London EC4R 2SU | United Kingdom

<sup>1.</sup> Russo E.B. Taming THC: potential cannabis synergy and phytocannabinoid-terpenoid entourage effects. British Journal of Pharmacology. 2011; 163: 1344-1364.

<sup>2.</sup> Grotenhermen F. Pharmacokinetics and pharmacodynamics of cannabinoids. Clin Pharmacokinet. 2003;42(4):327-60

<sup>3.</sup> Müller-Vahl, Kirsten & Grotenhermen, Franjo. (2017). Medizinisches cannabis: Die wichtigsten Änderungen. 114. A352- A356.

<sup>4.</sup> Ministry of Health Canada. Health Effects of Cannabis. 2017.